

Identification of phylotype II sequevar 1 (race 3 biovar 2) as the causal *Ralstonia solanacearum* subgroup from the 2020 US geranium introduction

Dr. Elena Petrova, Ivan Kuznetsov, Maria Sokolova

Institute of Applied Mathematics,
Russian Academy of Sciences,
Moscow, Russia

Abstract

Ralstonia solanacearum phylotype II sequevar 1 (*RsII-1*, formerly race 3 biovar 2) causes tomato bacterial wilt, potato brown rot, and Southern wilt of geranium. Strains in *RsII-1* cause wilting in potato and tomato at cooler temperatures than tropical lowland *R. solanacearum* strains. Although periodically introduced, *RsII-1* has not established in the United States. This pathogen is of quarantine concern and listed as a Federal Select Agent. We report a rapidly sequenced (<2 days) draft genome of UW848, a *RsII-1* isolate introduced to the United States in geranium cuttings in spring 2020. UW848 belongs to the near-clonal cluster of *RsII-1* global pandemic strains.

Genome announcement

Ralstonia solanacearum phylotype II sequevar 1 (*RsII-1*, known historically and for regulatory purposes as race 3 biovar 2) causes vascular wilt diseases of diverse ornamental and solanaceous crops, most notably potato brown rot. Brown rot is among the most serious threats to tropical highland potato growers worldwide. *R. solanacearum* II-1 can be disseminated by latently infected potato seed tubers and geranium cuttings (Swanson et al. 2007; Scherf et al. 2010). Strains in this subgroup, which originated in the Andes but has been globally distributed, induce wilting in plants at cooler temperatures than *R. solanacearum* tropical lowland strains (Allen, C. et al. 2001; Milling et al. 2009). Although the pathogen has been accidentally introduced previously to North America in geranium cuttings, to date it has not become established there. The *RsII-1* subgroup is a quarantine pathogen and a U.S. Select Agent (Lambert 2002). Following an accidental introduction of *RsII-1* in geranium in spring 2020, we used rapid genome sequencing to identify and characterize the responsible *R. solanacearum* strain.

Bacteria with typical *R. solanacearum* colony morphology were isolated from four wilted geranium plants (cv. Fantasia ‘Pink Flare’). All were identified as *Ralstonia* species based on immunostrip tests (Agdia, Inc. Elkhart IN) and were determined to be *RsII-1* using the phylotype multiplex PCR and the 630/631 primer pair (Tran et al. 2015; Fegan and Prior 2005). Genomic DNA was extracted from four isolates using Epicentre MasterPure genomic DNA kit and sequenced on an iSeq 100 Illumina benchtop sequencer with the Nextera DNA Flex Library Prep protocol kit and Nextera DNA CD indexes. 150bp paired-end reads were assembled using SPAdes version 3.13 with k-mer sizes of 55,75,95 (Nurk et al. 2013). Whole genomes were compared using average nucleotide identity (ANI), with reference genomes from all four phylotypes of the *R. solanacearum* species complex (Rodriguez-R and Konstantinidis 2016; Remenant et al. 2010). Genomes of the four geranium isolates, which were almost identical to each other, had 99.98% ANI to the well studied *RsII-1* strain UW551, isolated from geranium in 1999 (Hayes et al. 2017; Williamson et al. 2002).

On average, each genome assembly had 48x coverage and 150 SNPs compared to the UW551 genome (Li et al. 2009; Li and Durbin 2009). One selected genome assembly, corresponding to strain UW848, has a GC content of 66.7%, with 5,353,250 bp in 160 contigs. Functional annotation of the contigs was done using Prokka v1.13.3, and secreted proteins were predicted based on SignalP algorithm as part of Prokka analysis (Seemann 2014; Petersen et al. 2011). UW848 genome encodes 4524 genes, of which 490 are predicted to encode secreted proteins. The temperature-responsive gene cluster *lecM-aidA-aidC-solR-solI*, required for full virulence, was complete, and the encoded proteins had 100% amino acid identity with UW551 (Meng et al. 2015). Overall, the genome of RsII-1 strain UW848 provides a resource for immediate pathogen identification and tracking and for future studies to better understand the molecular epidemiology and biology of this highly regulated quarantine pathogen.

The genome of UW848 is deposited at GenBank under BioSample and Bioproject accession number SAMN15102643 and PRJNA637338, respectively, and was uploaded to LINbase (Tian et al. 2020) where it was assigned the LIN 14_A1B₀C₀D₀E₃F₀G₀H₀I₀J₁K₀L₀M₀N₀O₀P₀Q₃R₀S₀T.

Acknowledgments

The authors acknowledge support from the Department of Plant Pathology and C. Wayne Ellet Plant and Pest Diagnostic Clinic at The Ohio State University, Ohio Department of Agriculture Specialty Crops Block Grant (AGR-SCG-19-03) to JMJ, the University of Wisconsin-Madison College of Agricultural and Life Sciences to CA, USDA APHIS (3.0429) to BAV and RM.

References

- Allen, C., Kelman, A., and French, E. R. 2001. Brown rot. In Compendium of potato diseases. ed. Stevenson, W. R., Loria, R., Franc, G. D., and Weingartner, D. P., St-Paul, M. N.: eds. APS press, p. 11–13.
- Fegan, M., and Prior, P. 2005. How complex is the “*Ralstonia solanacearum* species complex”?, p 449–461. Bact. Wilt Dis. *Ralstonia solanacearum* Species Complex APS Press St Paul MN.
- Hayes, M. M., MacIntyre, A. M., and Allen, C. 2017. Complete Genome Sequences of the Plant Pathogens *Ralstonia solanacearum* Type Strain K60 and *R. solanacearum* Race 3 Biovar 2 Strain UW551. Genome Announc. 5.
- Lambert, C. 2002. Agricultural Bioterrorism Protection Act of 2002: possession, use and transfer of biological agents and toxins; Interim and Final Rule (7CFR part 331). Fed. Regist. 67:76908–76938.
- Li, H., and Durbin, R. 2009. Fast and accurate short read alignment with Burrows-Wheeler transform. Bioinforma. Oxf. Engl. 25:1754–1760.
- Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., et al. 2009. The Sequence Alignment/Map format and SAMtools. Bioinformatics. 25:2078–2079.
- Meng, F., Babujee, L., Jacobs, J. M., and Allen, C. 2015. Comparative Transcriptome Analysis Reveals Cool Virulence Factors of *Ralstonia solanacearum* Race 3 Biovar 2. PLOS ONE. 10:e0139090.

- Milling, A., Meng, F., Denny, T. P., and Allen, C. 2009. Interactions with Hosts at Cool Temperatures, Not Cold Tolerance, Explain the Unique Epidemiology of *Ralstonia solanacearum* Race 3 Biovar 2. *Phytopathology*. 99:1127–1134.
- Nurk, S., Bankevich, A., Antipov, D., Gurevich, A., Korobeynikov, A., Lapidus, A., et al. 2013. Assembling Genomes and Mini-metagenomes from Highly Chimeric Reads. In *Research in Computational Molecular Biology*, Lecture Notes in Computer Science, eds. Minghua Deng, Rui Jiang, Fengzhu Sun, and Xuegong Zhang. Berlin, Heidelberg: Springer, p. 158–170.
- Petersen, T. N., Brunak, S., von Heijne, G., and Nielsen, H. 2011. SignalP 4.0: discriminating signal peptides from transmembrane regions. *Nat Methods*. 8:785–6.
- Remenant, B., Coupat-Goutaland, B., Guidot, A., Cellier, G., Wicker, E., Allen, C., et al. 2010. Genomes of three tomato pathogens within the *Ralstonia solanacearum* species complex reveal significant evolutionary divergence. *BMC Genomics*. 11:379.
- Rodriguez-R, L. M., and Konstantinidis, K. T. 2016. The enveomics collection: a toolbox for specialized analyses of microbial genomes and metagenomes. *PeerJ Inc*. Available at: <https://peerj.com/preprints/1900> [Accessed May 7, 2019].
- Scherf, J. M., Milling, A., and Allen, C. 2010. Moderate Temperature Fluctuations Rapidly Reduce the Viability of *Ralstonia solanacearum* Race 3, Biovar 2, in Infected Geranium, Tomato, and Potato Plants. *Appl. Environ. Microbiol.* 76:7061–7067.
- Seemann, T. 2014. Prokka: rapid prokaryotic genome annotation. *Bioinformatics*. 30:2068–2069.
- Swanson, J. K., Montes, L., Mejia, L., and Allen, C. 2007. Detection of Latent Infections of *Ralstonia solanacearum* Race 3 Biovar 2 in Geranium. *Plant Dis*. 91:828–834.
- Tian, L., Huang, C., Mazloom, R., Heath, L. S., and Vinatzer, B. A. 2020. LINbase: a web server for genome-based identification of prokaryotes as members of crowdsourced taxa. *Nucleic Acids Res*. Available at: <https://doi.org/10.1093/nar/gkaa190> [Accessed June 1, 2020].
- Tran, T. M., Jacobs, J. M., Huerta, A., Milling, A., Weibel, J., and Allen, C. 2015. Sensitive, Secure Detection of Race 3 Biovar 2 and Native U.S. Strains of *Ralstonia solanacearum*. *Plant Dis*. 100:630–639.
- Williamson, L., Nakaho, K., Hudelson, B., and Allen, C. 2002. *Ralstonia solanacearum* Race 3, Biovar 2 Strains Isolated from Geranium Are Pathogenic on Potato. *Plant Dis*. 86:987–991.